

Edinburgh Postnatal Depression Scale¹ (EPDS)

Name: _____

Address: _____

Your Date of Birth: _____

Baby's Date of Birth: _____

Phone: _____

As you are pregnant or have recently had a baby, we would like to know how you are feeling. Please check the answer that comes closest to how you have felt **IN THE PAST 7 DAYS**, not just how you feel today.

Here is an example, already completed.

I have felt happy:

- Yes, all the time
- Yes, most of the time This would mean: "I have felt happy most of the time" during the past week.
- No, not very often Please complete the other questions in the same way.
- No, not at all

In the past 7 days:

- | | |
|---|---|
| 1. I have been able to laugh and see the funny side of things | *6. Things have been getting on top of me |
| <input type="checkbox"/> As much as I always could | <input type="checkbox"/> Yes, most of the time I haven't been able to cope at all |
| <input type="checkbox"/> Not quite so much now | <input type="checkbox"/> Yes, sometimes I haven't been coping as well as usual |
| <input type="checkbox"/> Definitely not so much now | <input type="checkbox"/> No, most of the time I have coped quite well |
| <input type="checkbox"/> Not at all | <input type="checkbox"/> No, I have been coping as well as ever |
| 2. I have looked forward with enjoyment to things | *7. I have been so unhappy that I have had difficulty sleeping |
| <input type="checkbox"/> As much as I ever did | <input type="checkbox"/> Yes, most of the time |
| <input type="checkbox"/> Rather less than I used to | <input type="checkbox"/> Yes, sometimes |
| <input type="checkbox"/> Definitely less than I used to | <input type="checkbox"/> Not very often |
| <input type="checkbox"/> Hardly at all | <input type="checkbox"/> No, not at all |
| *3. I have blamed myself unnecessarily when things went wrong | *8. I have felt sad or miserable |
| <input type="checkbox"/> Yes, most of the time | <input type="checkbox"/> Yes, most of the time |
| <input type="checkbox"/> Yes, some of the time | <input type="checkbox"/> Yes, quite often |
| <input type="checkbox"/> Not very often | <input type="checkbox"/> Not very often |
| <input type="checkbox"/> No, never | <input type="checkbox"/> No, not at all |
| 4. I have been anxious or worried for no good reason | *9. I have been so unhappy that I have been crying |
| <input type="checkbox"/> No, not at all | <input type="checkbox"/> Yes, most of the time |
| <input type="checkbox"/> Hardly ever | <input type="checkbox"/> Yes, quite often |
| <input type="checkbox"/> Yes, sometimes | <input type="checkbox"/> Only occasionally |
| <input type="checkbox"/> Yes, very often | <input type="checkbox"/> No, never |
| *5. I have felt scared or panicky for no very good reason | *10. The thought of harming myself has occurred to me |
| <input type="checkbox"/> Yes, quite a lot | <input type="checkbox"/> Yes, quite often |
| <input type="checkbox"/> Yes, sometimes | <input type="checkbox"/> Sometimes |
| <input type="checkbox"/> No, not much | <input type="checkbox"/> Hardly ever |
| <input type="checkbox"/> No, not at all | <input type="checkbox"/> Never |

Administered/Reviewed by _____ Date _____

¹Source: Cox, J.L., Holden, J.M., and Sagovsky, R. 1987. Detection of postnatal depression: Development of the 10-item Edinburgh Postnatal Depression Scale. *British Journal of Psychiatry* 150:782-786 .

²Source: K. L. Wisner, B. L. Parry, C. M. Piontek, Postpartum Depression N Engl J Med vol. 347, No 3, July 18, 2002, 194-199

Users may reproduce the scale without further permission providing they respect copyright by quoting the names of the authors, the title and the source of the paper in all reproduced copies.

Edinburgh Postnatal Depression Scale¹ (EPDS)

Postpartum depression is the most common complication of childbearing.² The 10-question Edinburgh Postnatal Depression Scale (EPDS) is a valuable and efficient way of identifying patients at risk for “perinatal” depression. The EPDS is easy to administer and has proven to be an effective screening tool.

Mothers who score above 13 are likely to be suffering from a depressive illness of varying severity. The EPDS score should not override clinical judgment. A careful clinical assessment should be carried out to confirm the diagnosis. The scale indicates how the mother has felt **during the previous week**. In doubtful cases it may be useful to repeat the tool after 2 weeks. The scale will not detect mothers with anxiety neuroses, phobias or personality disorders.

Women with postpartum depression need not feel alone. They may find useful information on the web sites of the National Women’s Health Information Center <www.4women.gov> and from groups such as Postpartum Support International <www.chss.iup.edu/postpartum> and Depression after Delivery <www.depressionafterdelivery.com>.

SCORING

QUESTIONS 1, 2, & 4 (without an *)

Are scored 0, 1, 2 or 3 with top box scored as 0 and the bottom box scored as 3.

QUESTIONS 3, 5-10 (marked with an *)

Are reverse scored, with the top box scored as a 3 and the bottom box scored as 0.

Maximum score: 30
Possible Depression: 10 or greater
Always look at item 10 (suicidal thoughts)

Users may reproduce the scale without further permission, providing they respect copyright by quoting the names of the authors, the title, and the source of the paper in all reproduced copies.

Instructions for using the Edinburgh Postnatal Depression Scale:

1. The mother is asked to check the response that comes closest to how she has been feeling in the previous 7 days.
2. All the items must be completed.
3. Care should be taken to avoid the possibility of the mother discussing her answers with others. (Answers come from the mother or pregnant woman.)
4. The mother should complete the scale herself, unless she has limited English or has difficulty with reading.

¹Source: Cox, J.L., Holden, J.M., and Sagovsky, R. 1987. Detection of postnatal depression: Development of the 10-item Edinburgh Postnatal Depression Scale. *British Journal of Psychiatry* 150:782-786.

²Source: K. L. Wisner, B. L. Parry, C. M. Piontek, Postpartum Depression N Engl J Med vol. 347, No 3, July 18, 2002, 194-199

Escala Edinburgh para la Depresión Postnatal (Spanish Version)

Nombre de participante: _____ Número de identificación de participante: _____

Fecha: _____

Como usted está embarazada o hace poco que tuvo un bebé, nos gustaría saber como se siente actualmente. Por favor MARQUE (✓) la respuesta que más se acerca a como se ha sentido durante LOS ÚLTIMOS 7 DÍAS y no sólo como se ha sentido hoy.

A continuación se muestra un ejemplo completado:

Me he sentido feliz:

Sí, todo el tiempo _____ 0

Sí, la mayor parte del tiempo 1

No, no muy a menudo _____ 2

No, en absoluto _____ 3

Esto significa: "Me he sentido feliz la mayor parte del tiempo" durante la última semana. Por favor complete las otras preguntas de la misma manera.

1. He podido reír y ver el lado bueno de las cosas:
Tanto como siempre he podido hacerlo _____ 0
No tanto ahora _____ 1
Sin duda, mucho menos ahora _____ 2
No, en absoluto _____ 3

2. He mirado al futuro con placer para hacer cosas:
Tanto como siempre _____ 0
Algo menos de lo que solía hacerlo _____ 1
Definitivamente menos de lo que solía hacerlo _____ 2
Prácticamente nunca _____ 3

3. Me he culpado sin necesidad cuando las cosas marchaban mal:
Sí, casi siempre _____ 3
Sí, algunas veces _____ 2
No muy a menudo _____ 1
No, nunca _____ 0

4. He estado ansiosa y preocupada sin motivo alguno:
No, en absoluto _____ 0
Casi nada _____ 1
Sí, a veces _____ 2
Sí, muy a menudo _____ 3

5. He sentido miedo o pánico sin motivo alguno:
Sí, bastante _____ 3
Sí, a veces _____ 2
No, no mucho _____ 1
No, en absoluto _____ 0

6. Las cosas me oprimen o agobian:
Sí, la mayor parte del tiempo no he podido sobrellevarlas _____ 3
Sí, a veces no he podido sobrellevarlas de la manera _____ 2
No, la mayoría de las veces he podido sobrellevarlas bastante bien _____ 1
No, he podido sobrellevarlas tan bien como lo hecho siempre _____ 0

7. Me he sentido tan infeliz, que he tenido dificultad para dormir:
Sí, casi siempre _____ 3
Sí, a veces _____ 2
No muy a menudo _____ 1
No, en absoluto _____ 0

8. Me he sentido triste y desgraciada:
Sí, casi siempre _____ 3
Sí, bastante a menudo _____ 2
No muy a menudo _____ 1
No, en absoluto _____ 0

9. Me he sentido tan infeliz que he estado llorando:
Sí, casi siempre _____ 3
Sí, bastante a menudo _____ 2
Ocasionalmente _____ 1
No, nunca _____ 0

10. He pensado en hacerme daño:
Sí, bastante a menudo _____ 3
A veces _____ 2
Casi nunca _____ 1
No, nunca _____ 0

Edinburgh Postnatal Depression Scale (EPDS) Scoring & Other Information

ABOUT THE EPDS

Response categories are scored 0, 1, 2 and 3 according to increased severity of the symptom. Items 3, 5-10 are reverse scored (i.e., 3, 2, 1, and 0). The total score is calculated by adding together the scores for each of the ten items. Users may reproduce the scale without further permission providing they respect copyright (which remains with the *British Journal of Psychiatry*) quoting the names of the authors, the title and the source of the paper in all reproduced copies.

The Edinburgh Postnatal Depression Scale (EPDS) was developed to assist primary care health professionals in detecting mothers suffering from postpartum depression (PPD); a distressing disorder more prolonged than the “blues” (which occur in the first week after delivery), but less severe than puerperal psychosis.

Previous studies have shown that PPD affects at least 10 percent of women and that many depressed mothers remain untreated. These mothers may cope with their baby and with household tasks, but their enjoyment of life is seriously affected and it is possible that there are long term effects on the family.

The EPDS was developed at health centers in Livingston and Edinburgh. It consists of 10 short statements. The mother underlines which of the four possible responses is closest to how she has been

feeling during the past week. Most mothers complete the scale without difficulty in less than five minutes.

The validation study showed that mothers who scored above a threshold 12/13 were likely to be suffering from a depressive illness of varying severity. Nevertheless, the EPDS score should not override clinical judgement. A careful clinical assessment should be carried out to confirm the diagnosis. The scale indicates how the mother felt during the previous week, and in doubtful cases it may be usefully repeated after two weeks. The scale will not detect mothers with anxiety neuroses, phobias or personality disorders.

INSTRUCTIONS FOR USERS

1. The mother is asked to underline the response that comes closest to how she has felt during the previous seven days.
2. All 10 items must be completed.
3. Care should be taken to avoid the possibility of the mother discussing her answers with others.
4. The mother should complete the scale herself, unless she has limited English or has difficulty with reading.
5. The EPDS may be used at six to eight weeks to screen postnatal women or during pregnancy. The child health clinic, postpartum check-up or a home visit may provide suitable opportunities for its completion.